arXiv:math/9906019v1 [math.NT] 2 Jun 1999

A characterization of the $\mathbf{Z}^n$ lattice

Noam D. Elkies

Introduction.

In this note we prove that $\mathbf{Z}^n$ is the only integral unimodular lattice $L \subset \mathbf{R}^n$ which does not contain a vector w such that $|w|^2 < n$ and $(v, v+w) \equiv 0 \bmod 2$ for all $v \in L$. By the work of Kronheimer and others on the Seiberg-Witten equation this yields an alternative proof of a theorem of Donaldson [D1,D2] on the geometry of 4-manifolds.

The proof uses the theory of theta series and modular forms; since this technique is not yet in the standard-issue arsenal of the 4-manifold community, I begin with an abbreviated exposition of this theory to make this note reasonably self-contained. This develops only the barest minimum, even to the point of never using the phrase "modular form"; for a more substantial exposition, refer to [Se, Ch.VII], and note the concluding remarks (6.7, "Complements").

Knowing that any $L \ncong \mathbf{Z}^n$ has characteristic vectors of norm $\leqslant n-8$, one might ask for which lattices is $n-8$ the minimum. It turns out that the same technique also yields a complete answer to this question. Since the answer may be of some interest (for instance there are 14 such lattices in each dimension $n \geqslant 23$), but its proof requires a somewhat more extensive use of modular forms, we announce the result at the end of this note but defer its proof and further discussion to a later paper.

Fractional linear transformations and theta series.

Let H be the Poincaré upper half-plane $\{t = x + iy : y > 0\}$, and let Γ be the group $\mathrm{PSL}_2(\mathbf{Z}) = \mathrm{SL}_2(\mathbf{Z})/\{\pm\mathbf{1}\}$, acting on H by fractional linear transformations:

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} : \; t \mapsto \frac{at+b}{ct+d} \; . \tag{1}$$

It is known that Γ is generated by $S = \left(\begin{smallmatrix} 0 & -1 \\ 1 & 0 \end{smallmatrix}\right)$ and $T = \left(\begin{smallmatrix} 1 & 1 \\ 0 & 1 \end{smallmatrix}\right)$, acting on H by

$$S(t) = -\frac{1}{t}, \quad T(t) = t+1. \tag{2}$$

Let $\Gamma(2) = \{\left(\begin{smallmatrix} a & b \\ c & d \end{smallmatrix}\right) \in \Gamma : b, c \text{ even}\}$; this is a normal subgroup of Γ, and reduction mod 2 yields the quotient map $\Gamma \to \Gamma/\Gamma(2) = \mathrm{PSL}_2(\mathbf{Z}/2) \cong \mathrm{S}_3$. Finally let $\Gamma_+ \subset \Gamma$ be the subgroup generated by S and T^2. Then Γ_+ has index 3 in Γ, contains $\Gamma(2)$ with index 2, and consists of the matrices congruent mod 2 to either $\mathbf{1}$ or S. Indeed it is clear that Γ_+ consists of matrices of this form; that all such matrices are in Γ_+ is perhaps most readily seen by proving as in [Se,

Ch.VII, Thm.1,2] that

$$D_+ := \{t = x + iy \in H : |x| \le 1, |t| \ge 1\} \tag{3}$$

(the ideal hyperbolic triangle in H with vertices $-1, 1, i\infty$) is a fundamental domain for the action of Γ_+ on H, and noting that D_+ is 3 times as large as the standard fundamental domain for Γ.

Now let L be a unimodular integral lattice in $\mathbf{R}^n$, i.e. a lattice of discriminant 1 such that $(v, v') \in \mathbf{Z}$ for all $v, v' \in L$. The theta series θ_L of L is a generating function encoding the norms $|v|^2 = (v, v)$ of lattice vectors:

$$\theta_L(t) := \sum_{v \in L} e^{\pi i |v|^2 t} \quad (t \in H). \tag{4}$$

For instance for $n = 1$ we have

$$\theta_{\mathbf{Z}}(t) := 1 + 2\left(e^{\pi i t} + e^{4\pi i t} + e^{9\pi i t} + \cdots\right) . \tag{5}$$

This sum converges uniformly in compact subsets of H (if $t = x + iy$ then $|e^{\pi i |v|^2 t}| = e^{-\pi |v|^2 y}$) and thus defines a holomorphic function on H. If L_1, L_2 are unimodular integral lattices in $\mathbf{R}^{n_1}, \mathbf{R}^{n_2}$ then $L_1 \oplus L_2$ is a unimodular integral lattice in $\mathbf{R}^{n_1+n_2}$ whose theta series is given by

$$\theta_{L_1 \oplus L_2}(t) = \theta_{L_1}(t) \cdot \theta_{L_2}(t). \tag{6}$$

Since each $|v|^2$ is an integer we have

$$\theta_L(t) = \theta_L(t + 2) = \theta_L(T^2(t)). \tag{7}$$

Since L is its own dual lattice we obtain a more interesting functional equation by applying Poisson inversion to (4):

$$(t/i)^{n/2} \theta_L(t) = \theta_L(-1/t) = \theta_L(S(t)), \tag{8}$$

where $(t/i)^{n/2}$ is the nth power of the principal branch of $\sqrt{t/i}$. By iterating (7,8) we find that for every $g = \left(\begin{smallmatrix} a & b \\ c & d \end{smallmatrix}\right)$ in $\langle S, T^2 \rangle = \Gamma_+$ there is a functional equation

$$\theta_L(g(t)) = \epsilon_n(c, d) \cdot (ct + d)^{n/2} \theta_L(t), \tag{9}$$

where again $(ct + d)^{n/2}$ is the nth power of the principal branch of $\sqrt{ct + d}$, and $\epsilon_n(c, d)$ is an eighth root of unity which does not depend on the choice of unimodular integral lattice L. (It does not depend on a, b because c, d determine g up to a power of T^2.) By choosing $L = \mathbf{Z}^n$ and using (6) we find

$$\epsilon_n(c, d) = \big(\epsilon_1(c, d)\big)^n. \tag{10}$$

Note that [Se, Ch.VII] assumes that L is an even lattice, i.e. $|v|^2 \in 2\mathbf{Z}$ for all $v \in L$. Such L have theta series invariant under T, and thus satisfy (9) for

all $g \in \langle S, T\rangle = \Gamma$. It is known from the arithmetic theory [Se, Ch.V] that $n \equiv 0 \bmod 8$ for such lattices, whence the ϵ_n factors all equal 1 in that case; this could also be proved analytically using (8) and the identity $(ST)^3 = \mathbf{1}$. We shall soon observe, en route to our estimate on the norm of characteristic vectors of odd lattices, that this method also yields an analytic proof of the fact [Se, Ch.V, Thm.2] that these vectors all have norm $\equiv n \bmod 8$.

How do fractional linear transformations $g \in \Gamma - \Gamma_+$ act on θ_L? We need only consider one representative of each of the two nontrivial cosets of Γ_+ in Γ, for instance $g = T$ and $g = TS$. For the first we find simply

$$\theta_L(T(t)) = \theta_L(t+1) = \sum_{v\in L} e^{\pi i|v|^2(t+1)} = \sum_{v\in L}(-1)^{|v|^2} e^{\pi i|v|^2 t}. \tag{11}$$

Now recall that the sign $v \mapsto (-1)^{|v|^2}$ is a group homomorphism from L to $\{\pm 1\}$ (because

$$|v+v'|^2 = |v|^2 + |v'|^2 + 2(v,v') \equiv |v|^2 + |v'|^2 \bmod 2 \tag{12}$$

for all $v, v' \in L$). Since L is unimodular there is a bijection between characters $L \to \{\pm 1\}$ and cosets of $2L$ in L which associates to the coset of any $w \in L$ the character $v \mapsto (-1)^{(v,w)}$. In particular there is a coset associated with $v \mapsto (-1)^{|v|^2}$; vectors in that coset, characterized by

$$|v|^2 \equiv (v,w) \bmod 2 \text{ for all } v \in L, \tag{13}$$

are known as <u>characteristic</u> <u>vectors</u> of L. (In [Se, Ch.V] this coset is called the "canonical class" in $L/2L$; in [CS2] this coset, scaled by $1/2$ to obtain a translate of L by $w/2$, is called the "shadow" of L, and our key formula (17) below is also a key ingredient of [CS2].) Choose some characteristic vector w, and rewrite (11) as

$$\theta_L(t+1) = \sum_{v\in L} e^{\pi i\left(|v|^2 t + (v,w)\right)}. \tag{14}$$

Applying Poisson inversion to this sum we find

$$(t/i)^{n/2}\theta_L(t+1) = \sum_{v\in L} e^{\pi i|v+\frac{w}{2}|^2(\frac{-1}{t})} = \theta'_L(S(t)), \tag{15}$$

where

$$\theta'_L(t) := \sum_{v\in L+\frac{w}{2}} e^{\pi i|v|^2 t} \tag{16}$$

is a generating function encoding the norms of canonical vectors. Replacing t by $St = -1/t$ in (16) we conclude that

$$\theta_L(TS(t)) = \theta_L(\frac{-1}{t}+1) = (t/i)^{n/2}\theta'_L(t). \tag{17}$$

To recover the result

$$|w|^2 \equiv n \bmod 8 \tag{18}$$

we may now regard (17) as a formula for $\theta_L'(t)$ and compare it with

$$\begin{aligned}\left(\frac{t+1}{i}\right)^{n/2}\theta_L'(t+1) &= \theta_L(TST(t)) = \theta_L(ST^{-1}S(t)) \\ = (T^{-1}S(t)/i)^{n/2}\theta_L(T^{-1}S(t)) &= \left(\frac{i(t+1)}{t}\right)^{n/2}\theta_L(TS(t))\end{aligned} \tag{19}$$

(in which we used $S^2 = (ST)^3 = \mathbf{1}$ and the invariance of θ_L under T^2, and again use $n/2$ power to mean nth power of principal square root). This yields

$$\theta_L'(t+1) = e^{\pi i n/4}\theta_L'(t). \tag{20}$$

Thus $\theta_L'(t)$ is a linear combination of terms $e^{\pi i m t/4}$ with $m \equiv n \bmod 8$, from which it follows that all the characteristic vectors have norm $\equiv n \bmod 8$ as claimed.

The characteristic vectors of $\mathbf{Z}$ are the odd integers, so

$$\theta_{\mathbf{Z}}'(t) = 2\sum_{m=0}^{\infty} e^{\pi i(m+\frac{1}{2})^2 t} = 2e^{\pi i t/4}\left(1 + e^{2\pi i t} + e^{6\pi i t} + e^{12\pi i t} + \cdots\right) . \tag{21}$$

Thus $\theta_{\mathbf{Z}}'(t) \sim 2e^{\pi i t/4} \to 0$ as $t \to i\infty$. From (17) it follows that $\theta_{\mathbf{Z}}$ tends to zero as $t \in D_+$ approaches the "cusp" ± 1. It will be crucial to us that $\theta_{\mathbf{Z}}$ **has no zeros in** H. This can be seen either from explicit product formulas such as

$$\sum_{m=0}^{\infty} q^{(m+\frac{1}{2})^2} = q^{1/4}\prod_{j=1}^{\infty}(1+q^{2j})(1-q^{4j}) \tag{22}$$

(a special case of the Jacobi triple product), or by using contour integrals as in [Se, Ch.VII, Thm.3] to show that ± 1 is the only zero of $\theta_{\mathbf{Z}}$ in $D_+ \cup \{\text{cusps}\}$. Also $\theta_{\mathbf{Z}}(i\infty) = 1$ so $\theta_{\mathbf{Z}}$ is bounded away from zero as $t \to i\infty$.

The shortest characteristic vector.

We are now ready to prove:

Theorem. *Let L be a unimodular integral lattice in $\mathbf{R}^n$ with no characteristic vector w such that $|w|^2 < n$. Then $L \cong \mathbf{Z}^n$.*

Proof: We first show that L and $\mathbf{Z}^n$ have the same theta function. To that end consider

$$R(t) := \theta_L(t)/\theta_{\mathbf{Z}^n}(t) = \theta_L(t)/\theta_{\mathbf{Z}}^n(t). \tag{23}$$

This is a holomorphic function because $\theta_{\mathbf{Z}}$ does not vanish in H. Since θ_L and $\theta_{\mathbf{Z}^n}$ both transform according to (9) under Γ_+, their quotient $R(t)$ is invariant under Γ_+. By the hypothesis on L we have $\theta_L' \ll e^{\pi i n t/4}$ as $t \to i\infty$. Thus $\theta_L'/\theta_{\mathbf{Z}^n}'$ is bounded as $t \to i\infty$, whence by (17) $R(t)$ is bounded as $t \in D_+$ approaches ± 1. Finally $R(i\infty) = 1$. By the maximum principle we deduce that R is the constant function 1, i.e. $\theta_L = \theta_{\mathbf{Z}}^n$.

Thus for each m the lattices L and $\mathbf{Z}^n$ have the same number of vectors of norm m. Taking $m = 1$ we find that L has n pairs of unit vectors. Since L is integral these must be orthogonal to each other, and thus generate a copy of $\mathbf{Z}^n$ inside L. Using integrality again we conclude that this copy is all of L. QED

Since the hypothesis is automatically satisfied if $n < 8$ we also recover the fact that $\mathbf{Z}^n$ is the only unimodular integral lattice for those n. With some more work we can also use the relation between θ_L and θ'_L and the theory of modular forms to completely describe those $L \subset \mathbf{R}^n$ whose shortest characteristic vector has norm $n - 8$: these are precisely the lattices of the form $\mathbf{Z}^{n-r} \oplus L_0$, where $L_0 \subset \mathbf{R}^r$ is a unimodular integral lattice with no vectors of norm 1 and exactly $2n(23 - n)$ vectors of norm 2. In particular $n \leqslant 23$ and there are only finitely many choices for L_0. Fortunately the table of unimodular lattices in [CS1, pp.416–7] extends just far enough that we can list all possible L_0. These are tabulated below, indexed as in the table of [CS1] by the root system of norm-2 vectors:

r	8	12	14	15	16	17	18	18	19	20	20	21	22	23
	E_8	D_{12}	E_7^2	A_{15}	D_8^2	$A_{11}E_6$	D_6^3	A_9^2	$A_7^2D_5$	D_4^5	A_5^4	A_3^7	A_1^{22}	O_{23}

Of these the first is the E_8 lattice and the last is the "shorter Leech lattice"; these are the unimodular integral lattices of minimal dimension having minimal norm 2 and 3 respectively. It also follows from the analysis that each of these lattices has exactly $2^{n-11}r$ characteristic vectors of norm $n - 8$. We defer the proof of the $\mathbf{Z}^{n-r} \oplus L_0$ criterion, and an analogous condition for self-dual binary codes, to a subsequent paper.

Acknowledgements. Thanks to Tom Mrowka for bringing this problem to my attention and to John H. Conway for enlightening correspondence. This work was made possible in part by funding from the National Science Foundation and the Packard Foundation.

References

[Do1] Donaldson, S.K.: An application of gauge theory to the topology of 4-manifolds. *J. Diff. Geom.* **18** (1983), 279–315.

[Do2] Donaldson, S.K.: The orientation of Yang-Mills moduli spaces and 4-manifold topology. *J. Diff. Geom.* **26** (1987), 397–428.

[CS1] Conway, J.H., Sloane, N.J.A.: *Sphere Packings, Lattices and Groups.* New York: Springer 1993.

[CS2] Conway, J.H., Sloane, N.J.A.: A new upper bound on the minimal distance of self-dual codes. *IEEE Trans. Inform. Theory* **36** (1990), 1319–1333.

[Se] Serre, J.-P.: *A Course in Arithmetic.* New York: Springer, 1973.

Dept. of Mathematics
Harvard University
Cambridge, MA 02138 USA
e-mail: `elkies@math.harvard.edu`

January, 1995